

Pharmacy



Medical Necessity Criteria for Phospodiesterase-5 Inhibitors – Cialis (tadalafil), Levitra (vardenafil), Staxyn (vardenafil ODT)

Background - After evaluating the relative clinical and cost effectiveness of the PDE-5 inhibitors for erectile dysfunction, the DoD P&T Committee recommended that Cialis (tadalafil) and Levitra (vardenafil) and Staxyn (vardenafil ODT) be designated as non-formulary. This recommendation has been approved by the Director, TMA. Prior authorization requirements and quantity limits continue to apply to the PDE-5 inhibitors.

Effective Date: 18 April 2012

Medical Necessity Criteria for Cialis, Levitra and Staxyn

The non-formulary cost share for Cialis, Levitra and Staxyn may be reduced to the formulary cost share IF one or more of the following criteria are met:

- 1. The patient had a hypersensitivity reaction to Viagra
- 2. The patient has experienced significant adverse effects from Viagra
- 3. **For a patient previously treated with Cialis, Levitra or Staxyn:** the patient tried Viagra for at least 90 days, titrated to the maximum recommended dose, and experienced significant decrease in erectile function compared to previous therapy.
- 4. **For a patient not previously treated with Cialis, Levitra or Staxyn:** the patient tried Viagra for at least 90 days, titrated to the maximum recommended dose, and experienced no improvement in erectile function.
- 5. The patient is being treated with Cialis for pulmonary arterial hypertension (previously referred to as primary pulmonary hypertension).

Criteria approved through the DoD P&T Committee process November 2011

www.tricare.mil is the official Web site of the TRICARE Management Activity, a component of the <u>Military Health System</u> Skyline 5, Suite 810, 5111 Leesburg Pike, Falls Church, VA 22041-3206



TRICARE Pharmacy Program Medical Necessity Form for Cialis (tadalafil), Levitra (vardenafil), and Staxyn (vardenafil)



has resulted in therapeutic failure.

Step 3

565B

This form applies to the TRICARE Pharmacy Program (TPharm). The medical necessity criteria outlined on this form also apply at Military Treatment Facilities (MTFs). The form must be completed and signed by the prescriber.

- Viagra is the formulary agent of this drug class on the DoD Uniform Formulary at the formulary cost share. Cialis, Levitra, and Staxyn are nonformulary, but available to most beneficiaries at the non-formulary cost share
- You do NOT need to complete this form in order for non-Active duty beneficiaries (spouses, dependents, and retirees) to obtain non-formulary medications at the non-formulary cost share. The purpose of this form is to provide information that will be used to determine if the use of a non-formulary medication is medically necessary. If a non-formulary medication is determined to be medically necessary, non-Active duty beneficiaries may obtain it at the formulary cost share.
- Active duty service members may not fill prescriptions for a non-formulary medication unless it is determined to be medically necessary. There is no

MAIL ORDER and And RETAIL	The proof the control of the control of the part to the part to the part of t	The provider may call: 1-866-684-4488 or the completed form may be faxed to: 1-866-684-4477 The patient may attach the completed for to the prescription and mail it to: Expres P.O. Box 52150, Phoenix, AZ 85072-99 or email the form only to: TPharmPA@express-scripts.com		M M	Non-formulary medications are available at MTFs only if both of the following are met: The prescription is written by a military provider or, at the discretion of the MTF, a civilian provider to whom the patient was referred by the MTF. The non-formulary medication is determined to be medically necessary. Please contact your local MTF for more information. There are no cost shares at MTFs.	
Step Plea	se complete	patient and physici	ian information	(please	e print):	
Patient Name:					Physician Name:	
Addr	ess:				Address:	
Sponsor ID #					Phone #:	
Date of Birth:					Secure Fax #:	
					Secure rax #.	
You M unacc					ary medication. Circle a reason use of the formulary medication	
You M unacc	UST supply a eptable.	specific written clinic			ary medication. Circle a reas y use of the formulary medic	
You M unacc	UST supply a eptable. Formulary edication a (sildenafil)	specific written clinic	cal explanation a	s to wh	ary medication. Circle a reas y use of the formulary medic Clinical Explanation	
You Munacc	eptable. Formulary edication a (sildenafil) stable clinical se of the formu	Reason 1 2 3 4 5 6 7 reasons for not using lary medication is contri	the formulary maindicated.	s to why	ary medication. Circle a reason use of the formulary medication Clinical Explanation on(s) are:	
You M unacc F M Viagr Accep 1. Us 2. Th	eptable. Formulary edication a (sildenafil) stable clinical se of the formulary edication	Reason 1 2 3 4 5 6 7 reasons for not using lary medication is contrexperienced significant	the formulary maindicated. adverse effects fr	edication	ry medication. Circle a reason use of the formulary medication Clinical Explanation on(s) are:	
You Munacc FM Viagr Accep 1. Us 2. Th 3. Us	eptable. Formulary edication a (sildenafil) Stable clinical se of the formulary edication	Reason 1 2 3 4 5 6 7 reasons for not using lary medication is contrexperienced significant lary medication has resulting the second contract of the second con	the formulary mandicated. adverse effects from the sulted in the same and the sulted in the same and the same	edication om the fittic failure	ry medication. Circle a reason use of the formulary medication Clinical Explanation on(s) are:	cation would be

The patient is being treated for the preservation/restoration of erectile function following prostatectomy AND use of Viagra

Date

The patient is being treated with Cialis for signs and symptoms of BPH (benign prostatic hypertrophy).

The patient is being treated with Cialis for pulmonary arterial hypertension (PAH).

Prescriber Signature

I certify the above is true to the best of my knowledge. Please sign and date: